

3-1976

Surgical Techniques for Arthrodesis of Two to Four Adjacent Spinal Vertebrae Throughout the Entire Spinal Column

Kent K. Wu

Follow this and additional works at: <https://scholarlycommons.henryford.com/hfhmedjournal>



Part of the [Life Sciences Commons](#), [Medical Specialties Commons](#), and the [Public Health Commons](#)

Recommended Citation

Wu, Kent K. (1976) "Surgical Techniques for Arthrodesis of Two to Four Adjacent Spinal Vertebrae Throughout the Entire Spinal Column," *Henry Ford Hospital Medical Journal* : Vol. 24 : No. 1 , 39-50.

Available at: <https://scholarlycommons.henryford.com/hfhmedjournal/vol24/iss1/7>

This Article is brought to you for free and open access by Henry Ford Health System Scholarly Commons. It has been accepted for inclusion in Henry Ford Hospital Medical Journal by an authorized editor of Henry Ford Health System Scholarly Commons.

Surgical Techniques for Arthrodesis of Two to Four Adjacent Spinal Vertebrae Throughout the Entire Spinal Column

Kent K. Wu, MD*

Surgical techniques for cervical, thoracic and lumbar spinal fusions have been described. So far, 51 such fusions have been performed with satisfactory results.

Spinal fusion can be defined as a surgical procedure by which a surgeon attempts to produce synostosis of two or more adjacent spinal vertebrae. Generally, its three main objectives are:

1. To provide stability of the spine.
2. To correct or prevent deformity of the spine.
3. To relieve pain and suffering from diseases of the spine.

Based on clinical experience, diseases and deformities which may require spinal fusion include the following:

1. Trauma—fractures and dislocations.
2. Congenital anomalies—hemivertebra, scoliosis, spondylolisthesis, etc.
3. Degenerative diseases—osteoarthritis, ruptured disks, etc.
4. Inflammatory diseases—rheumatoid arthritis, ankylosing spondylitis, etc.
5. Infections—caused by bacteria, fungi, parasites, etc.
6. Tumors—benign, malignant or metastatic tumors that cause significant destruction of the spine.
7. Metabolic bone diseases and osteoporosis, osteomalacia, etc.
8. Endocrine disorders—hyperparathyroidism, hyperthyroidism, hyperpituitarism, etc.
9. Diseases of skeletal muscle—different myopathies, neuromyopathies, etc.
10. Hereditary diseases—Marfan's syndrome, osteogenesis imperfecta, Ehlers-Danlos syndrome, etc.
11. Diseases of the nervous system—polio-myelitis, cerebral palsy, meningo-myelocoele, etc.

* Department of Orthopedic Surgery

This is the first in a series of papers by Dr. Wu which will describe techniques and experience at Henry Ford Hospital

Address reprint requests to Dr. Wu at Department of Orthopedic Surgery, Henry Ford Hospital, 2799 West Grand Boulevard, Detroit, MI 48202

From an anatomical point of view, five different surfaces are available for spinal fusion on each vertebra. They include the vertebral body, the two transverse processes, the two superior and inferior articular processes, the two laminae and the spinous process. However, because the anterior surface of the transverse processes and the vertebral body are situated in the anterior aspect of the vertebral column, it naturally follows that there are two general approaches to the spine, namely, the anterior and the posterior approaches.

In surgical operations, either an anterior or posterior spinal fusion can be done. The anterior approach is used to gain access to the anterior surface of the transverse process and the vertebral bodies. The posterior approach enables the surgeon to expose the posterior surface of the transverse processes, the articular processes, the laminae and the spinous processes. The anterior spinal fusion is commonly called an interbody fusion. Posterior spinal fusions can be posteromedial or posterolateral. The posteromedial fusion usually includes the spinous processes, the laminae and the articular processes. The posterolateral fusion normally incorporates only the transverse processes and the articular processes. Due to the numerous indications for spinal fusion, many surgical techniques and their results have been published in the medical literature.¹⁻¹¹⁶

The surgical techniques described in this paper are the ones we routinely use.

Techniques

The "H" graft technique for C1-C2 fusion (Figures 1-2)

An incision about 4" in length is made along the midline of the posterior aspect of the neck extending from the external occipital protuberance to the spinous process of C3. The cervical paraspinal muscles are stripped from the posterior arch of atlas and the spinous process and laminae of axis.




Figure 1
The stainless steel wire has been passed around the iliac graft and the posterior elements of C1 and C2 vertebrae prior to its tightening.




Figure 2
The stainless steel wire has been tightened and the iliac graft is now firmly wired to the posterior arches of atlas and axis.

These exposed bone surfaces are roughened with a sharp bone rasp. Next, a notch about 1/4" in depth is made at the superior base of C2 spinous process with a small rongeurs and a small hole is also made through the same spinous process slightly above the notch with a sharp right-angled awl. Using a small rongeurs, four notches are cut in the four sides of the rectangular iliac graft with the top and bottom notches along the midline and the two lateral notches more proximally based. A size 20 stainless steel wire loop is passed upward under the posterior arch of atlas and the sculptured iliac graft is placed on the posterior arch of atlas, care

Arthrodesis of the Spine

being taken to firmly engage the inferior notch of the graft with the superior notch of C2 spinous process. The two ends of the wire are made to engage the two lateral notches of the iliac graft and are passed through the wire loop. Tension is applied to the two ends of the wire simultaneously. As the slack is taken out, the iliac graft is forced to make intimate contact with the posterior arch of atlas. The two ends of the wire are passed under tension through the hole at the base of C2 spinous process in opposite directions. They are then turned downward and are tightly twisted together against the inferior base of C2 spinous process. The incision is closed in the usual manner.

The dual onlay iliac graft technique for fusing two to four spinal vertebrae from C-2 to T-12 (Figures 3-7)

After skin incision and stripping of paraspinal muscles at the appropriate spinal level, an air-powered bur is used to decorticate partially the spinous processes, the laminae and the posterior aspect of the adjacent articular processes. Small holes are made through the bases of the spinous processes with a sharp right-angled awl. Two pieces of mirror-image iliac grafts are obtained from the outer table of ilium. Holes matching the holes of the spinous processes in size and distance are drilled through the iliac grafts with a small bur. With size 20 stainless steel wire, the two grafts are tightly wired to the decorticated spinous processes and the laminae of the spinal segments to be fused. Match-stick iliac grafts are placed over the decorticated articular processes. The incision is closed in the regular manner.

Although the dual onlay iliac graft technique can be used in lumbar spinal fusion, our experience shows that very large pieces of iliac bone are required to accomplish the fusion due to the relatively large dimensions of the lumbar vertebrae. Consequently, we prefer the spinous process splitting technique for our lumbar spinal fusion.




Figure 3
A transparent view of the posterior aspect of C5, C6 and C7 vertebrae.




Figure 4
The cervical paraspinal muscles have been stripped from the posterior arches of C5, C6 and C7 vertebrae.




Figure 5

The dual iliac grafts are about to be wired to the decorticated spinous process and laminae.




Figure 6

The dual iliac grafts have been firmly wired in place.

The spinous process splitting technique for lumbar spinal fusion (Figures 8-13)

After the skin incision and stripping of lumbar paraspinal muscles have been accomplished, the basal portions of the spinous processes, the laminae and the articular processes are decorticated with a sharp spinal gouge. The supraspinous ligaments, the interspinous ligaments and the spinous processes of the vertebra to be fused are split down the midline almost to the neural arches with a stryker saw and a wide

osteotome. A properly prepared iliac graft is forcefully inserted into the split space bridging the spinous processes together. Holes are made through the bases of the spinous processes and size 20 stainless steel wires are used to hold the iliac graft tightly between the split spinous processes. Match-stick iliac grafts are placed over the decorticated spinous processes, laminae and articular processes. The incision is closed in the usual manner. Sometimes a lumbar spinal fusion is needed following a wide posterior decompression lumbar laminectomy which usually removes the spinous processes, the laminae and parts of the articular processes. In these circumstances, the surgeon can perform a posterolateral fusion which can be accomplished by partially decorticating the adjacent articular processes and the transverse processes with an air-powered burr. This is followed by placing long and slender iliac grafts in the gutter formed by the articular processes and the transverse processes of the involved vertebrae (Figures 14-15).

Postoperative managements

The cervical fusion is usually protected with a plaster minerva cast for about three months followed by the use of a four poster cervical brace or a cervical collar for an additional two to three months.

Arthrodesis of the Spine




Figure 7

The iliac bone chips have been placed over the decorticated articular processes.




Figure 8

A transparent view of the posterior aspect of the lumbosacral spine.




Figure 9

The lumbar paraspinal muscles have been stripped from the posterior arches of L3, L4 and L5 vertebrae.




Figure 10

The spinous processes are being split with a stryker saw and the posterior arches are being decorticated with a spinal gouge.




Figure 11

The iliac graft is being inserted into the split spinous processes and the laminae and the articular processes can also be decorticated with an air-powered burr.

Arthrodesis of the Spine




Figure 12
The iliac graft is being wired to the split spinous processes.




Figure 13
Iliac bone chips have been placed over the decorticated laminae and articular processes.




Figure 14
The articular processes and the transverse processes are being decorticated with an air-powered burr.




Figure 15
Long and slender iliac bone grafts have been placed in the gutter formed by the decorticated articular processes and the transverse processes.

Thoracic and lumbar fusions are managed in a plaster body cast for about three months. Then a back brace is worn for an additional two to three months.

About 24 hours after the cast is applied, the patient is urged to get up and ambulate for the purpose of preventing postoperative pulmonary, vascular, urinary and skin complications. After discharge from the hospital,

patients are seen on an outpatient basis at monthly intervals. They are also encouraged to pursue light daily activities.

Discussion

Our spinal fusion techniques have been utilized in 51 spinal fusions. They include three C1-C2 fusions, three lower cervical fusions, three thoracic fusions and 42 lumbar and lumbosacral fusions. So far, our results have been exceptionally good. A detailed analysis of all our spinal fusion cases will be published when a much bigger series has been accumulated.

Summary

Surgical techniques for fusing two to four spinal segments throughout the entire spinal column have been described. So far, 51 such fusions have been performed with very encouraging results.

Acknowledgement

Thanks to Mr. Jay Knipstein for his superb drawings.

References

1. Adkins EWO: Lumbo-sacral arthrodesis after laminectomy. *J Bone Joint Surg* **37B**:208, 1955
2. Albee FH: A report of bone transplantation and osteoplasty in the treatment of Pott's disease of spine. *NY J Med* **95**:469, 1912
3. Albee FH: Transplantation of a portion of the tibia into the spine for Pott's disease. A preliminary report. *JAMA* **57**:885, 1911
4. Allison N: Fusion of the spinal column. *Surg Gynec Obstet* **46**:826, 1928
5. Aronson N, Filtzer DL, and Bagan M: Anterior cervical fusion by the Smith-Robinson approach. *J Neurosurg* **29**:397, 1968
6. Bailey RW, and Badgley CE: Stabilization of cervical spine by anterior fusion, *J Bone Joint Surg* **42A**:565, 1960
7. Baker LD, and Hoyt WA, Jr.: The use of interfacet Vitallium screws in the Hibbs fusion, *Southern Med J* **41**:419, 1948
8. Benton BF, and Calandruccio RA: Surgical technic of anterior lumbar fusion. *Amer Surg* **32**:134, 1966
9. Bosworth DM: Circumduction fusion of the spine. *J Bone Joint Surg* **38A**:263, 1956
10. Bosworth DM: Clothespin graft of the spine for spondylolisthesis and laminal defects. *Amer J Surg* **67**:61, 1945

Arthrodesis of the Spine

11. Bosworth DM: Surgery of the spine. American Academy of Orthopaedic Surgeons Instructional Course Lectures, vol. 14, Ann Arbor, 1957, JW Edwards
12. Bosworth DM: Techniques of spinal fusion: pseudarthrosis and method of repair, American Academy of Orthopaedic Surgeons Instructional Course Lectures, vol. 5, Ann Arbor, 1948, JW Edwards
13. Bosworth DM: Technique of spinal fusion in the lumbosacral region by the double clothespin graft (distraction graft: "H" graft) and results, American Academy of Orthopaedic Surgeons Instructional Course Lectures, vol. 9, Ann Arbor, 1952, JW Edwards
14. Bosworth DM, Wright HA, Fielding JW, and Goodrich ER: A study in the use of bank bone for spine fusion in tuberculosis. *J Bone Joint Surg* **35A**:329, 1953
15. Boucher HH: A method of spinal fusion. *J Bone Joint Surg* **41B**:248, 1959
16. Calandruccio RA, and Benton BF: Anterior lumbar fusion. In DePalma, AF, editor: *Clinical Orthopaedics and Related Research*, vol 35, Philadelphia, 1964, JB Lippincott Co
17. Campbell WC: Operative measures in the treatment of low back pain. *Trans Southern Surg Assoc* **42**:207, 1929
18. Campbell WC: Operative measures in the treatment of affections of the lumbosacral and sacroiliac articulation. *Surg Gynec Obstet* **51**:381, 1930
19. Cattell HS, and Clark GL Jr.: Cervical kyphosis and instability following multiple laminectomies in children. *J Bone Joint Surg* **49A**:713, 1967
20. Cleveland M: Technique of spine fusion for tuberculosis involving vertebra. American Academy of Orthopaedic Surgeons Instructional Course Lectures, vol 9, Ann Arbor, 1952, JW Edwards
21. Cleveland M, Bosworth DM, Fielding JW and Smyrnis P: Fusion of the spine for tuberculosis in children. A long-range follow-up study. *J Bone Joint Surg* **40A**:91, 1958
22. Cloward RB: Lesions of the intervertebral disks and their treatment by interbody fusion methods. The painful disk. In DePalma, AF, editor: *Clinical Orthopaedics and Related Research*, vol 27, Philadelphia, 1963, JB Lippincott Co
23. Cloward RB: New method of diagnosis and treatment of cervical disc disease. *Clin Neurosurg* **8**: 93, 1962
24. Cloward RB: The treatment of ruptured lumbar intervertebral discs by vertebral body fusion. I. Indications, operative technique, after care, *J Neurosurg* **10**:154, 1953
25. Cloward RB: Vertebral body fusion for ruptured cervical discs. Description of instruments and operative technic. *Amer J Surg* **98**:722, 1959
26. Cobb JR: Technique, after-treatment, and results of spine fusion for scoliosis. American Academy of Orthopaedic Surgeons Instructional Course Lectures, vol 9, Ann Arbor, 1952, JW Edwards
27. Connor AC, Rooney JA, and Carroll JP: Anterior lumbar fusion. A technique combining intervertebral and intravertebral body fixation. *Surg Clin N Amer* **47**:231, 1967
28. Cook WA, Shaw RR, Webb WR, Clark WK, and Shah HH: Transthoracic evacuation and anterior spine fusion in Pott's paraplegia. *Ann Thorac Surg* **4**:291, 1967
29. Curran JP, and McGaw WH: Posterolateral spinal fusion with pedicle grafts. In Urist MR, editor: *Clinical Orthopaedics and Related Research*, vol 59, Philadelphia, 1968, JB Lippincott Co
30. DePalma AF, and Cooke AJ: Results of anterior interbody fusion of the cervical spine. *Clin Orthop* **60**:169, 1968
31. DePalma AF, and Prabhakar M: Posterior-posterobilateral fusion of the lumbosacral spine. In Urist MR, editor: *Clinical Orthopaedics and Related Research*, vol 47, Philadelphia, 1966, JB Lippincott Co
32. DePalma AF, and Ray RD: Surgery of the lumbar spine. *Clin Orthop* **63**:162, 1969
33. Dommisse GF: Lumbo-sacral interbody spinal fusion. *J Bone Joint Surg* **41B**:87, 1959
34. Fang HSY, Ong GB, and Hodgson AR: Anterior spinal fusion, the operative approaches. In DePalma AF, editor: *Clinical Orthopaedics and Related Research*, vol 35, Philadelphia, 1964, JB Lippincott Co
35. Fielding JW, Lusskin R, and Batista A: Multiple segment anterior cervical spinal fusion. In Urist MR, editor: *Clinical Orthopaedics and Related Research*, vol 54, Philadelphia, 1967, JB Lippincott Co

Kent K. Wu

36. Gallie WE: Fractures and dislocations of the cervical spine. *Amer J Surg* **46**:495, 1939
37. Garber JN: Surgery in cervical spine injuries. American Academy of Orthopaedic Surgeons Instructional Course Lectures, vol 18, St. Louis, 1961, The CV Mosby Co, p 67
38. Garrett AL, Perry J, and Nickel VL: Stabilization of the collapsing spine. *J Bone Joint Surg* **43A**:474, 1961
39. Gibson A: A modified technique for spinal fusion. *Surg Gynec Obstet* **53**:365, 1931
40. Goldstein LA: Results of treatment of scoliosis with turnbuckle plaster cast correction and fusion, *J Bone Joint Surg* **41A**:321, 1959.
41. Goldstein LA: The surgical management of scoliosis. *Clin Orthop* **35**:95, 1964
42. Goldstein LA: Surgical management of scoliosis. *J Bone Joint Surg* **48A**:167, 1966
43. Goldstein LA: The surgical management of scoliosis. *Clin Orthop* **77**:32, 1971
44. Goldstein LA: The surgical treatment of idiopathic scoliosis. Part I. Selection of the fusion area. *Clin Orthop* **93**:131, 1973
45. Goldstein LA: The surgical treatment of idiopathic scoliosis. Part II. Indications, technique, results. *Clin Orthop* **93**:137, 1973
46. Goldstein LA: Treatment of idiopathic scoliosis by Harrington instrumentation and fusion with fresh autogenous iliac bone grafts. Results in 80 patients, *J Bone Joint Surg* **51A**:209, 1969
47. Goldstein LA, and Evarts CM: Further experiences with the treatment of scoliosis by cast correction and spine fusion with fresh autogenous iliac bone grafts, *J Bone Joint Surg* **48A**:962, 1966
48. Handra BE: Wiring the spinous processes in Pott's disease. *Trans Amer Orthop Assoc* **4**:206, 1891
49. Hallock H, Francis KC, and Jones JB: Spine fusion in young children. A long-term end-result study with particular reference to growth effects. *J Bone Joint Surg* **39A**:481, 1957
50. Hallock H, and Jones JB: Tuberculosis of the spine. An end-result study of the effects of the spine fusion operation in a large number of patients *J Bone Joint Surg* **36A**:219, 1954
51. Harmon PH: Anterior excision and vertebral body fusion operation for intervertebral disk syndromes of the lower lumbar spine: Three-to-five-year results in 244 cases. In DePalma, AF, editor: *Clinical Orthopaedics and Related Research*, vol 26, Philadelphia, 1963, JB Lippincott Co
52. Harmon PH: Anterior extraperitoneal lumbar disc excision and vertebral body fusion. In DePalma, AF, editor: *Clinical Orthopaedics*, vol 18, Philadelphia, 1960, JB Lippincott Co
53. Harmon PH: Subtotal anterior lumbar disc excision and vertebral body fusion. III. Application to complicated and recurrent multilevel degenerations. *Amer J Surg* **97**:649, 1959
54. Harrington PR: Treatment of scoliosis. Correction and internal fixation by spine instrumentation. *J Bone Joint Surg* **44A**:591, 1962
55. Henderson ED: Results of the surgical treatment of spondylolisthesis, *J Bone Joint Surg* **48A**:619, 1966
56. Henderson MS: Operative fusion for tuberculosis of the spine. *JAMA* **92**:45, 1929
57. Henry MO, and Geist ES: Spinal fusion by simplified technique, *J Bone Joint Surg* **15**:622, 1933
58. Hibbs RA: A further consideration of an operation for Pott's disease of the spine, *Ann Surg* **55**:682, 1912
59. Hibbs RA: An operation for progressive spinal deformities, *NY Med J* **93**:1013, 1911
60. Hibbs RA and Risser JC: Treatment of vertebral tuberculosis by the spine fusion operation. A report of 286 cases. *J Bone Joint Surg* **10**:805, 1928
61. Hibbs RA, Risser JC, and Ferguson AB: Scoliosis treated by the fusion operation. An end result study of 360 cases. *J Bone Joint Surg* **13**:91, 1931
62. Hodgson, AR: An approach to the cervical spine C3-C7. *Clin Orthop* **39**:129, 1965

Arthrodesis of the Spine

64. Hodgson AR and Wong SK: A description of a technic and evaluation of results in anterior spinal fusion for deranged intervertebral disk and spondylolisthesis. In Urist MR, editor: *Clinical Orthopaedics and Related Research*, vol 56, Philadelphia, 1968, JB Lippincott Co
65. Holdsworth FW: Fractures, dislocations, and fracture-dislocations of the spine. *J Bone Joint Surg* **45B**:6, 1963
66. Hoover NW: Methods of lumbar fusion. *J Bone Joint Surg* **50A**:194, 1968
67. Howorth B: Low backache and sciatica: Results of surgical treatment. Part I. Spine fusion only. *J Bone Joint Surg* **46A**:1485, 1964
68. Howorth B: Low backache and sciatica: Results of surgical treatment. Part II. Removal of nucleus pulposus and spine fusion. *J Bone Joint Surg* **46A**:1500, 1964
69. Howorth B: Low backache and sciatica: Results of surgical treatment. Part III. Surgical treatment of spondylolisthesis. *J Bone Joint Surg* **46A**:1515, 1964
70. Howorth MB: Evolution of spinal fusion. *Ann Surg* **117**:278, 1943
71. Jaslow IA: Intercorporeal bone graft in spinal fusion after disc removal. *Surg Gynec Obstet* **82**:215, 1946
72. Johnson JT, and Robinson RA: Anterior strut grafts for severe kyphosis. Results of 3 cases with a preceding progressive paraplegia. In Urist MR, editor: *Clinical Orthopaedics and Related Research*, vol 56, Philadelphia, 1968, JB Lippincott Co
73. King D: Internal fixation for lumbosacral fusion. *Amer J Surg* **66**:357, 1944
74. Kite JH: Nonoperative versus operative treatment of tuberculosis of the spine in children; review of 50 consecutive cases treated by each method. *Southern Med J* **26**:918, 1933
75. Lange F: The operative splinting of the vertebral column in Pott's disease. *Surg Gynec Obstet* **44**:668, 1927
76. Law WA: Osteotomy of the spine. *J Bone Joint Surg* **44A**:1199, 1962
77. McBride ED, and Shorbe HB: Lumbosacral fusion: the mortised transfacet method by use of the vibrating electric saw for circular bone blocks. In DePalma AF, editor: *Clinical Orthopaedics*, vol 12, Philadelphia, 1958, JB Lippincott Co
78. McMaster PE: Osteotomy of the spine for fixed flexion deformity. *J Bone Joint Surg* **44A**:1207, 1962
79. Mercer W: Spondylolisthesis: With a description of a new method of operative treatment and notes of ten cases. *Edinburgh Med J* **43**:545, 1936
80. Moe JH: A critical analysis of methods of fusion for scoliosis. An evaluation in 266 patients. *J Bone Joint Surg* **40A**:529, 1958
81. Overton LM.: An improved technic for arthrodesis of the lumbosacral spine. *Amer J Surg* **80**:559, 1950
82. Overton LM: Arthrodesis of the lumbosacral spine. In DePalma AF, editor: *Clinical Orthopaedics*, vol 5, Philadelphia, 1955, JB Lippincott Co
83. Overton, LM: Lumbosacral arthrodesis: an evaluation of its present status. *Amer Surg* **25**:771, 1959
84. Pennal GF, McDonald GA, and Dale GG: A method of spinal fusion using internal fixation. In DePalma AF, editor: *Clinical Orthopaedics and Related Research*, vol 35, Philadelphia, 1964, JB Lippincott Co
85. Perry J, and Nickel V L: Total cervical spine fusion for neck paralysis. *J Bone Joint Surg* **41A**:37, 1959
86. Petter CK: Rib-splinter graft in spinal fusion for vertebral tuberculosis. *J Bone Joint Surg* **19**:413, 1937
87. Robinson RA: Anterior and posterior cervical spine fusions. *Clin Orthop* **35**:34, 1964
88. Robinson RA: Anterior and posterior cervical spine fusions. In DePalma AF, editor: *Clinical Orthopaedics and Related Research*, vol 35, Philadelphia, 1964, JB Lippincott Co
89. Robinson RA, and Southwick WO: Surgical approaches to the cervical spine. American Academy of Orthopaedic Surgeons Instructional Course Lectures, vol 17, St. Louis, 1960, The CV Mosby Co, p 299

90. Robinson RA, Walker EA, Ferlic DC, and Wiecking DK: The results of anterior interbody fusion of the cervical spine. *J Bone Joint Surg* **44A**:1569, 1962
91. Rombald C.: Treatment of spondylolisthesis by postero-lateral fusion, resection of the pars interarticularis, and prompt mobilization of the patient. *J Bone Joint Surg* **48A**:1282, 1966
92. Ryerson EW: Surgical treatment of low back disabilities. *J Bone Joint Surg* **14**:154, 1932
93. Sacks S: Anterior interbody fusion of the lumbar spine. *J Bone Joint Surg* **47B**:211, 1965
94. Sacks S: Anterior interbody fusion of the lumbar spine. Indications and results in 200 cases. In DePalma AF, editor: *Clinical Orthopaedics and Related Research*, vol 44, Philadelphia, 1966, JB Lippincott Co
95. Simmons EH, and Bhalla SK: Anterior cervical discectomy and fusion. A clinical and biomechanical study with eight-year follow-up. *J Bone Joint Surg* **51B**:225, 1969
96. Smith A DeF: Lumbosacral fusion by the Hibbs technique. American Academy of Orthopaedic Surgeons Instructional Course Lectures, vol 9, Ann Arbor, 1952, JW Edwards
97. Smith A DeF: Tuberculosis of the spine. Results in 70 cases treated at the New York Orthopaedic Hospital from 1945 to 1960. In Urist MR, editor: *Clinical Orthopaedics and Related Research*, vol 58, Philadelphia, 1968, JB Lippincott Co
98. Smith GW, and Robinson RA: The treatment of certain cervical spine disorders by anterior removal of the intervertebral disc and interbody fusion. *J Bone and Joint Surg* **40A**:607, 1958
99. Smith-Petersen MN, Larson CB, and Aufranc OE: The classic: Osteotomy of the spine for correction of flexion deformity in rheumatoid arthritis, *Clin Orthop* **66**:6, 1969
100. Southwick WO, and Robinson RA: Recent advances in surgery of the cervical spine. *Surg Clin N Amer* **41**:1661, 1961
101. Speed K: Spondylolisthesis: Treatment by anterior bone graft. *Arch Surg* **37**:175, 1938
102. Stinchfield FE, and Sinton WA: Criteria for spine fusion with use of "H" bone graft following disc removal. Results in 100 cases, *Arch Surg* **65**:542, 1952
103. Stock FE: Anterior spinal fusion. A review of five years' work, *Aust New Zeal J Surg* **31**:161, 1962
104. Tambornino JM, Armbrust EN, and Moe JH: Harrington instrumentation in correction of scoliosis. A comparison with cast correction. *J Bone Joint Surg* **46A**:313, 1964
105. Toumey JW: Internal fixation in fusion of the lumbosacral joint. *Lahey Clin Bull* **3**:188, 1943
106. Truchly and Thompson, WAL: Posterolateral fusion of the lumbosacral spine. *J Bone Joint Surg* **44A**:505, 1962
107. Verbiest H: Anterolateral operations for fractures and dislocations in the middle and lower parts of the cervical spine. Report of a series of forty-seven cases, *J Bone Joint Surg* **51A**:1489, 1969
108. Watkins MB: Posterolateral fusion of the lumbar and lumbosacral spine. *J Bone Joint Surg* **35A**:1014, 1953
109. Watkins MB: Posterolateral bone-grafting for fusion of the lumbar and lumbosacral spine. *J Bone Joint Surg* **41A**:388, 1959
110. Watkins MB: Posterolateral fusion in pseudarthrosis and posterior element defects of the lumbosacral spine. In DePalma AF, editor: *Clinical Orthopaedics and Related Research*, vol 35, Philadelphia, 1964, JB Lippincott Co
111. Williams JL, Allen MB Jr, and Harkess JW: Late results of cervical discectomy and interbody fusion: Some factors influencing the results. *J Bone Joint Surg* **50A**:277, 1968
112. Wilson PD, and Straub LR: Lumbosacral fusion with metallic-plate fixation. American Academy of Orthopaedic Surgeons Instructional Course Lectures, vol 9, Ann Arbor, 1952, JW Edwards
113. Wiltberger BR: The dowel intervertebral-body fusion as used in lumbar-disc surgery, *J Bone Joint Surg* **39A**:284, 1957
114. Wiltberger BR: Intervertebral body fusion by the use of posterior bone dowel. In DePalma AR, editor: *Clinical Orthopaedics and Related Research*, vol 35, Philadelphia, 1964, JB Lippincott Co
115. Wiltse LL: Spondylolisthesis in children. In DePalma AF, editor: *Clinical Orthopaedics* vol 21, Philadelphia, 1961, JB Lippincott Co
116. Young HH, and Love JG: End results of removal of protruded lumbar intervertebral discs with and without fusion. American Academy of Orthopaedic Surgeons Instructional Course Lectures vol. 16, Ann Arbor, 1959, JW Edwards, p 213